


Parental Smoking and Childhood Brain Tumors

A population-based case-control study of
the role of functional polymorphisms

Jessica Trimis, MA, MS, Roberta McKean-Cowdin, PhD,
Susan Searles Nielsen, PhD, Beth Mueller DrPH

University of Southern California, Keck School of Medicine, Department of Preventive Medicine
Fred Hutchinson Cancer Research Center, Seattle, Washington

Parental smoking during pregnancy and risk of CBT



Boffetta, 2000; Huncharek, 2002

Methods

- **Subset of population based case-control study (202 Cases / 286 Controls)**
- **Smoking exposure (in-person interview):**
 - Maternal: never/ever and by dose
 - Paternal: never/ever and duration of exposure
- **Genotyping—archival newborn blood spots**
 - 9 functional polymorphisms in 6 genes
- **OR and 95% CI by logistic regression**
 - Stratification and formal interaction tests

Table 1. Parental smoking during pregnancy and risk of CBT


Exposure	Cases N(%) N=149	Controls N(%) N=210	Adj. OR (95% CI)
No exposure to prenatal smoke	104 (69.8)	153 (72.9)	1.00
Father smoked during pregnancy	45 (30.2)	57 (27.1)	1.03 (0.62, 1.71)
Father only	25 (16.8)	27 (12.9)	1.24 (0.66, 2.35)
Father and other source (mother or other resident)	20 (13.4)	30 (14.3)	0.82 (0.41, 1.63)

Exposure	Cases N(%) N = 125	Controls N(%) N = 200	Adj. OR (95% CI)
No exposure to prenatal smoke	104 (83.2)	153 (76.5)	1.00
Mother smoked during pregnancy	21 (16.8)	47 (23.5)	0.55 (0.29, 1.05)
Mother only	4 (3.2)	12 (6.0)	0.41 (0.12, 1.42)
Mother and other source (father or other resident)	17 (13.6)	35 (17.5)	0.60 (0.30, 1.21)

Table 2. Parental smoking, EPHX1 genotype and risk of CBT

Exposure	Normal PAH Activity		High PAH Activity		p-value for interaction	
	Cases/Controls	Adj. OR (95%CI)	Cases/Controls	Adj. OR (95%CI)		
<i>Paternal Smoking</i>						
EPHX1 H139R	Never	107/144	1.00	44/82	1.00	0.07
	<3.5 hours	12/18	0.74 (0.33, 1.65)	12/15	1.32 (0.52, 3.34)	
	3.5+ hours	12/19	0.96 (0.42, 2.20)	9/5	3.18 (0.92, 10.98)	
<i>p for trend</i>			0.71		0.07	
<i>Maternal Smoking</i>						
Exposure	Normal PAH Activity		High PAH Activity		p-value for interaction	
	Cases/Controls	Adj. OR (95%CI)	Cases/Controls	Adj. OR (95%CI)		
<i>Maternal Smoking</i>						
EPHX1 H139R	Never	123/149	1.00	52/87	1.00	0.003
	1-10/day	3/18	0.17 (0.05, 0.62)	2/8	0.28 (0.05, 1.50)	
	11+/day	5/14	0.46 (0.15, 1.41)	11/7	2.19 (0.72, 6.63)	
<i>p for trend</i>			0.02		0.38	

Exposure to paternal smoking and risk of CBT by genotype for EPHX1 H139R



Conclusion

- Modest association between paternal smoking and CBT risk
- CBT risk associated with paternal smoking is greater in children with high activity EPHX1 genotype
- Environmental risk factors for CBT may be better detected when considering genetic background

Acknowledgements

- Additionally, we would like to thank the following people for their support in this project: Dr. Susan Preston Martin, Dr. Elizabeth Holly, Dr. Federico M. Farin, and the Washington State Department of Health Newborn Screening Program (Michael Glass and Michael Ginder), California Department of Public Health Genetic Disease Screening Program (Steve Graham, Marty Kharrazi, and Fred Lorey), the Sequoia Foundation for obtaining specimens, and the Functional Genomics Core Laboratory, Center for Ecogenetics and Environmental Health, University of Washington (Jesse Tsai and Hannah-Malia A. Viernes) for genotyping

Parental smoking and risk of CBT by genotype

