

PON1 polymorphisms and Wilms Tumor


Gisele Moledo de Vasconcelos, PhD

**Gonçalves BAA, Azevedo RM, Pombo-de-Oliveira MS, de Camargo B and the
Brazilian Embryonic Tumor Group**


Pediatric Hematology-Oncology Program

Instituto Nacional de Câncer, Rio de Janeiro, Brazil


Wilms tumor




- Leukemias
- Neuroblastoma
- Soft tissues
- Lymphomas
- Other
- Kidney
- CNS
- Bones



Fetal kidney
18 weeks



Wilms tumor



Mature kidney

EXPOSITION TO ENVIRONMENTAL RISK FACTORS



Father

Mother

Mother/Child

Mother

Child

Pre-conceptual

During Pregnancy

Post-natal

Pesticides and Wilms Tumor

- Previous studies have shown an association of pesticides exposures and Wilms Tumor risk.

British Journal of Cancer (1998) 77(5), 825–829

© 1998 Cancer Research Campaign

Childhood cancer and paternal employment in agriculture: the role of pesticides

NT Fear¹, E Roman², G Reeves¹ and B Pannett³

¹Cancer Epidemiology Unit, Imperial Cancer Research Fund, Gibson Building, Radcliffe Infirmary, Woodstock Road, Oxford, OX2 6HE; ²Leukaemia Research Fund, Centre for Clinical Epidemiology, University of Leeds, Leeds, LS2 9JT; ³Medical Research Council Environmental Epidemiology Unit, University of Southampton, Southampton General Hospital, Southampton, SO16 6YD, UK



ELSEVIER

Available online at www.sciencedirect.com



Int. J. Hyg. Environ.-Health 209 (2006) 57–64

**International Journal
of Hygiene and
Environmental Health**

www.elsevier.de/ijheh

Research | Children's Health

Wilms' tumor and exposures to residential and occupational hazardous chemicals

James Tsai^{*}, Wendy E. Kaye, Frank J. Bove

Centers for Disease Control and Prevention (CDC), Agency for Toxic Substances and Disease Registry (ATSDR),
1600 Clifton Road, Mailstop-E86, Atlanta, GA 30333, USA

Household Pesticides and the Risk of Wilms Tumor

Maureen A. Cooney,¹ Julie L. Daniels,¹ Julie A. Ross,² Norman E. Breslow,³ Brad H. Pollock,⁴ and Andrew F. Olshan¹

¹Department of Epidemiology, University of North Carolina, Chapel Hill, North Carolina, USA; ²Department of Pediatrics, University of Minnesota, Minneapolis, Minnesota, USA; ³Department of Biostatistics, University of Washington, Seattle, Washington, USA; ⁴Department of Epidemiology and Biostatistics, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

Pesticides and Wilms Tumor



American Journal of Epidemiology

Copyright © 1995 by The Johns Hopkins University School of Hygiene and Public Health

All rights reserved

Vol. 141, 1


Printed in U.S.A.

Parental Exposures to Pesticides and Risk of Wilms' Tumor in Brazil

Colin R. Sharpe,¹ Eduardo L. Franco,¹ Beatriz de Camargo,² L. Fernando Lopes,² J. Henrique Barreto,³
Rosanne R. Johnsson,³ and Marcos A. Maud³

Paraoxonase 1 (PON1)

- Hydrolyses active metabolites of organophosphorus pesticides.
- Levels of PON1 change according to age.




- And also due to polymorphisms.

Polymorphisms in the gene *PON1*

PON1 Q192R – change activity

PON1 L55M – change enzyme levels



Children susceptibility to Wilms Tumor can be modified by *PON1*

genotypes:

their own genotype


and during pregnancy, in case of exposition, maternal genotype

Aims


- Identify the distribution of *PON1* (Q192R and L55M) polymorphisms and its relationship with Wilms tumor & embryonal neuroectodermal origin tumors development in Brazilian children.
- Identify the frequencies of *PON1* polymorphisms in mothers and the occurrence of embryonic tumors in their children.

Subjects and Methods

- Allelic discrimination (TaqMAN probes)



- 47 Brazilian children with WT and their mothers;
- 49 other embryonal tumors (retinoblastoma, neuroblastoma and medulloblastoma) and their mothers;
- 190 health children and 117 mothers



Subjects and Methods

- Wilms Tumor patients: age ranged from 0 – 125 months.
- Other embryonal tumors: age range from 0 – 132 months.
- Children used as controls: age range from 0 – 60 months.
- There was no statistical difference according to gender in tumors nor even in controls.

Results

Table II: Genotype distributions and polymorphisms susceptibility

		Wilms Tumor		Neuroectodermal Tumors	
Genotype		Child	Mothers	Child	Mothers
		OR (95%CI)	OR (95%CI)	OR (95%CI)	OR (95%CI)
<i>PONI Q192R</i>	QQ	1.0	1.0	1.0	1.0
	QR	2.25 (0.98 - 5.14)	1.58 (0.68 - 3.66)	1.61 (0.78 - 3.30)	1.04 (0.49 - 2.19)
	RR	1.83 (0.71 - 4.7)	1.64 (0.56 - 4.81)	1.47 (0.64 - 3.34)	2 (0.81 - 4.92)
<i>PONI L55M</i>	LL	1.0	1.0	1.0	1.0
	LM	1.72 (0.87 - 3.41)	1.45 (0.68 - 3.12)	0.91 (0.48 - 1.74)	0.69 (0.35 - 1.37)
	MM	1.12 (0.34 - 3.62)	2.28 (0.58 - 8.86)	0.33 (0.07 - 1.49)	0.43 (0.08 - 2.36)

- Mothers genotypes and children genotypes together:
- *PON1* Q192R – QQ/QQ; QQ/QR; QR/QQ; QR/QR; RR/QR; RR/RR
- *PON* L55M – LL/LL; LL/LM; LM/LL; LM/MM; MM/LM; MM/MM
- We observed no significant differences but N is too small.

Final Remark

- Despite the small series of cases , our preliminary results suggest that *PON1* Q192R polymorphism may be an important risk factor in the development of WT when children are exposed to pesticides.

Acknowledgements

- Ministério da Saúde
- Support Agencies
- Pediatric Hematology-Oncology Program
- Brazilian Embryonic tumor group
- Dra. Maria do Socorro Pombo-de-Oliveira
& Dra. Beatriz de Camargo
- Rafaela Montalvão & Bruno A. Gonçalves



Ministério da
Saúde



THANK YOU!

